

REC'D 20 MAY 2003

WIPO

PCT

P1 1009827

# THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE  
United States Patent and Trademark Office

May 15, 2003

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM  
THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK  
OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT  
APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A  
FILING DATE.

APPLICATION NUMBER: 60/371,946

FILING DATE: April 11, 2002

RELATED PCT APPLICATION NUMBER: PCT/US03/11297



By Authority of the  
COMMISSIONER OF PATENTS AND TRADEMARKS

P. SWAIN  
Certifying Officer

**PRIORITY DOCUMENT**  
SUBMITTED OR TRANSMITTED IN  
COMPLIANCE WITH  
RULE 17.1(a) OR (b)

04/11/02  
JCS65 U.S. TO

"EXPRESS MAIL CERTIFICATE"

"Express Mail" Mailing Label Number EL737872380US

Date Of Deposit: April 11, 2002

I Hereby Certify That This Paper Or Fee Is Being Deposited With The United States Postal Service "Express Mail Post Office To Addressee" Service Under 37 CFR 1.10 On The Date Indicated Above And Is Addressed To: Assistant Commissioner for Patents, Washington, D.C. 20231.

Name Of Person Mailing Paper Or Fee,

(Type Or Print) Elsa Matos

Signature Elsa Matos

04-12-02

A/P102

U.S. PRO  
60/371946  
04/11/02

PROVISIONAL APPLICATION COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION for PATENT under 37 CFR 1.53(c).

Docket No.		P51337P	
INVENTOR(s) / APPLICANT(s)			
Last Name	First Name	Middle Initial	Residence (City and Either State or Foreign Country)
CALLAHAN	James	F.	King of Prussia, Pennsylvania 19406
LI	Yue	H.	Collegeville, Pennsylvania 19426

TITLE OF THE INVENTION (280 characters max)			
NF-KB INHIBITORS			
Correspondence Address:			
GLAXOSMITHKLINE			
Corporate Intellectual Property - UW2220		Telephone No. 610-270-5019	
709 Swedeland Road		Facsimile No. 610-270-5090	
King of Prussia			
State	PA	Zip Code	19406-0939
Country	United States of America		

ENCLOSED APPLICATION PARTS (check all that apply)			
<input checked="" type="checkbox"/> Specification	Number of Pages	26	<input type="checkbox"/> Small Entity Statement
<input checked="" type="checkbox"/> Abstract	Number of Pages	1	
<input type="checkbox"/> Drawings	Number of Sheets		<input type="checkbox"/> Other (specify)

METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT		
<input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge filing fees and credit Deposit Account No. 19-2570	PROVISIONAL FILING FEE AMOUNT (\$)	\$160.00

Respectfully submitted,  
Signature:

Soma G. Simon  
Soma G. Simon

Date:

4/11/02

Registration No.: 37,444

☐ Additional inventors are being named on separately numbered sheets attached hereto.

PROVISIONAL APPLICATION FILING ONLY

SEND TO: Assistant Commissioner for Patents, Box Provisional Application, Washington, D.C. 20231.



20462

PATENT TRADEMARK OFFICE

N:\SOMA\CASES\IP51337P\ProvApp.doc

**NF- $\kappa$ B INHIBITORS****FIELD OF THE INVENTION**

- This invention relates in general to a method of inhibiting pathological activation of the transcription factor NF- $\kappa$ B (nuclear factor- $\kappa$ B) using
- 5 aminothiophene compounds. Such methods are particularly useful for treating diseases in which activation of NF- $\kappa$ B is implicated. More specifically, these methods may be used for inhibiting IKK- $\beta$  (I $\kappa$ B kinase- $\beta$ ) phosphorylation of I $\kappa$ B (inhibitory protein  $\kappa$ B)-which prevents subsequent degradation and activation of NF- $\kappa$ B dimers. Such methods are useful in the treatment of a variety of diseases
- 10 associated with NF- $\kappa$ B activation including inflammatory and tissue repair disorders; particularly rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease) osteoarthritis; osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and ultraviolet radiation (UV)-induced skin damage; autoimmune diseases including systemic lupus
- 15 erythematosus, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherosclerosis, restenosis, diabetes, glomerulonephritis, cancer, including Hodgkins disease, cachexia, inflammation associated with infection and certain viral infections, including acquired immune deficiency syndrome (AIDS), adult respiratory distress syndrome, Ataxia
- 20 Telangiectasia.

**BACKGROUND OF THE INVENTION**

- Recent advances in scientific understanding of the mediators involved in acute and chronic inflammatory diseases and cancer have led to new strategies in the search for effective therapeutics. Traditional approaches include direct target
- 25 intervention such as the use of specific antibodies, receptor antagonists, or enzyme inhibitors. Recent breakthroughs in the elucidation of regulatory mechanisms involved in the transcription and translation of a variety of mediators have led to increased interest in therapeutic approaches directed at the level of gene transcription.

Nuclear factor  $\kappa$ B (NF- $\kappa$ B) belongs to a family of closely related dimeric transcription factor complexes composed of various combinations of the Rel/NF- $\kappa$ B family of polypeptides. The family consists of five individual gene products in mammals, RelA (p65), NF- $\kappa$ B1 (p50/p105), NF- $\kappa$ B2 (p49/p100), c-Rel, and RelB, all of which can form hetero- or homodimers. These proteins share a highly homologous 300 amino acid "Rel homology domain" which contains the DNA binding and dimerization domains. At the extreme C-terminus of the Rel homology domain is a nuclear translocation sequence important in the transport of NF- $\kappa$ B from the cytoplasm to the nucleus. In addition, p65 and cRel possess potent transactivation domains at their C-terminal ends.

The activity of NF- $\kappa$ B is regulated by its interaction with a member of the inhibitor I $\kappa$ B family of proteins. This interaction effectively blocks the nuclear localization sequence on the NF- $\kappa$ B proteins, thus preventing migration of the dimer to the nucleus. A wide variety of stimuli activate NF- $\kappa$ B through what are likely to be multiple signal transduction pathways. Included are bacterial products (LPS), some viruses (HIV-1, HTLV-1), inflammatory cytokines (TNF $\alpha$ , IL-1), environmental and oxidative stress and DNA damaging agents. Apparently common to all stimuli however, is the phosphorylation and subsequent degradation of I $\kappa$ B. I $\kappa$ B is phosphorylated on two N-terminal serines by the recently identified I $\kappa$ B kinases (IKK- $\alpha$  and IKK- $\beta$ ). Site-directed mutagenesis studies indicate that these phosphorylations are critical for the subsequent activation of NF- $\kappa$ B in that once phosphorylated the protein is flagged for degradation via the ubiquitin-proteasome pathway. Free from I $\kappa$ B, the active NF- $\kappa$ B complexes are able to translocate to the nucleus where they bind in a selective manner to preferred gene-specific enhancer sequences. Included in the genes regulated by NF- $\kappa$ B are a number of cytokines and chemokines, cell adhesion molecules, acute phase proteins, immunoregulatory proteins, eicosanoid metabolizing enzymes and anti-apoptotic genes.

It is well-known that NF- $\kappa$ B plays a key role in the regulated expression of a large number of pro-inflammatory mediators including cytokines such as TNF, IL-1 $\beta$ , IL-6 and IL-8, cell adhesion molecules, such as ICAM and VCAM, and

inducible nitric oxide synthase (iNOS). Such mediators are known to play a role in the recruitment of leukocytes at sites of inflammation and in the case of iNOS, may lead to organ destruction in some inflammatory and autoimmune diseases.

The importance of NF- $\kappa$ B in inflammatory disorders is further strengthened by studies of airway inflammation including asthma, in which NF- $\kappa$ B has been shown to be activated. This activation may underlie the increased cytokine production and leukocyte infiltration characteristic of these disorders. In addition, inhaled steroids are known to reduce airway hyperresponsiveness and suppress the inflammatory response in asthmatic airways. In light of the recent findings with regard to glucocorticoid inhibition of NF- $\kappa$ B, one may speculate that these effects are mediated through an inhibition of NF- $\kappa$ B.

Further evidence for a role of NF- $\kappa$ B in inflammatory disorders comes from studies of rheumatoid synovium. Although NF- $\kappa$ B is normally present as an inactive cytoplasmic complex, recent immunohistochemical studies have indicated that NF- $\kappa$ B is present in the nuclei, and hence active, in the cells comprising rheumatoid synovium. Furthermore, NF- $\kappa$ B has been shown to be activated in human synovial cells in response to stimulation with TNF- $\alpha$  or IL-1 $\beta$ . Such a distribution may be the underlying mechanism for the increased cytokine and eicosanoid production characteristic of this tissue. See Roshak, A. K., et al., *J. Biol. Chem.*, **271**, 31496-31501 (1996). Expression of IKK- $\beta$  has been shown in synoviocytes of rheumatoid arthritis patients and gene transfer studies have demonstrated the central role of IKK- $\beta$  in stimulated inflammatory mediator production in these cells. See Aupperle et al. *J. Immunology* 1999. 163:427-433 and Aupperle et al. *J. Immunology* 2001;166:2705-11. More recently, the intra-articular administration of a wild type IKK- $\beta$  adenoviral construct was shown to cause paw swelling while intra-articular administration of dominant-negative IKK- $\beta$  inhibited adjuvant-induced arthritis in rat. See Tak et al. *Arthritis and Rheumatism* 2001; 44:1897-1907.

The NF- $\kappa$ B/Rel and I $\kappa$ B proteins are also likely to play a key role in neoplastic transformation and metastasis. Family members are associated with cell transformation *in vitro* and *in vivo* as a result of overexpression, gene amplification,

gene rearrangements or translocations. In addition, rearrangement and/or amplification of the genes encoding these proteins are seen in 20-25% of certain human lymphoid tumors. Further, NF- $\kappa$ B is activated by oncogenic ras, the most common defect in human tumors and blockade of NF- $\kappa$ B activation inhibits ras mediated cell transformation. In addition, a role for NF- $\kappa$ B in the regulation of apoptosis has been reported, strengthening the role of this transcription factor in the regulation of tumor cell proliferation. TNF, ionizing radiation and DNA damaging agents have all been shown to activate NF- $\kappa$ B which in turn leads to the upregulated expression of several anti-apoptotic proteins. Conversely, inhibition of NF- $\kappa$ B has been shown to enhance apoptotic-killing by these agents in several tumor cell types. As this likely represents a major mechanism of tumor cell resistance to chemotherapy, inhibitors of NF- $\kappa$ B activation may be useful chemotherapeutic agents as either single agents or adjunct therapy. Recent reports have implicated NF- $\kappa$ B as an inhibitor of skeletal cell differentiation as well as a regulator of cytokine-induced muscle wasting (Guttridge et al. *Science*; 2000; 289: 2363-2365.) further supporting the potential of NF- $\kappa$ B inhibitors as novel cancer therapies.

Several NF- $\kappa$ B inhibitors are described in C. Wahl, et al. *J. Clin. Invest.* 101(5), 1163-1174 (1998), R. W. Sullivan, et al. *J. Med. Chem.* 41, 413-419 (1998), J. W. Pierce, et al. *J. Biol. Chem.* 272, 21096-21103 (1997)

The marine natural product hymenialdisine is known to inhibit NF- $\kappa$ B. Roshak, A., et al., *JPET*, 283, 955-961 (1997). Breton, J. J and Chabot-Fletcher, M. C., *JPET*, 282, 459-466 (1997).

Additionally, patent applications have been filed on indole and benzimidazole inhibitors of the IKK complex, see DE 19928424 and WO200130774, and the natural products staurosporine, quercetin, K252a and K252b have been shown to be IKK- $\beta$  inhibitors, see Peet, G. W. and Li, J. J. *Biol. Chem.*, 274, 32655-32661 (1999) and Wisniewski, D., et al., *Analytical Biochem.* 274, 220-228 (1999).

U.S. Patent No. 3,963,750 describes the preparation of certain aminothiophenes.

### SUMMARY OF THE INVENTION

The present invention involves novel compounds and novel methods of inhibiting the activation transcription factor NF- $\kappa$ B using the present compounds.

5        An object of the present invention is to provide a method for treating diseases which may be therapeutically modified by altering the activity of transcription factor NF- $\kappa$ B.

Accordingly, in the first aspect, this invention provides a pharmaceutical composition comprising a compound according to Formula I.


10        In another aspect, this invention provides a method of treating diseases in which the disease pathology may be therapeutically modified by inhibiting phosphorylation and subsequent degradation of I $\kappa$ B by IKK- $\beta$ .

In still another aspect, this invention provides a method of treating diseases in which the disease pathology may be therapeutically modified by inhibiting  
15        pathological activation of NF- $\kappa$ B.

In a particular aspect, this invention provides methods for treating a variety of diseases associated with NF- $\kappa$ B activation including inflammatory and tissue repair disorders, particularly rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease) osteoarthritis,  
20        osteoporosis and fibrotic diseases, dermatosis, including psoriasis, atopic dermatitis and ultraviolet radiation (UV)-induced skin damage; autoimmune diseases including systemic lupus erythematosus, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherosclerosis, restenosis, diabetes, glomerulonephritis, cancer, including Hodgkins disease,  
25        cachexia, inflammation associated with infection and certain viral infections, including acquired immune deficiency syndrome (AIDS), adult respiratory distress syndrome and Ataxia Telangiectasia

**DETAILED DESCRIPTION OF THE INVENTION**

The compounds of the present invention are selected from Formula (I) herein below:



(I)

wherein:

- R<sub>1</sub> represents NR<sub>4</sub>R<sub>5</sub>;
- 10 R<sub>2</sub> represents CONH<sub>2</sub> or SO<sub>2</sub>NH<sub>2</sub>;
- R<sub>3</sub> is selected from the group consisting of halogen, C<sub>1-4</sub>alkyl, NH<sub>2</sub>, CF<sub>3</sub>, OCF<sub>3</sub>, O-alkyl, S-alkyl, CN, CHO, SO<sub>2</sub>-alkyl, (CH<sub>2</sub>)<sub>q</sub>NR<sub>7</sub>R<sub>8</sub>, O-(CH<sub>2</sub>)<sub>q</sub>NR<sub>7</sub>R<sub>8</sub> and NO<sub>2</sub>;
- R<sub>4</sub> represents C<sub>1-4</sub>alkyl;
- R<sub>5</sub> represents CONHR<sub>6</sub>;
- 15 R<sub>6</sub> is selected from the group consisting of hydrogen, alkyl and aryl;
- R<sub>7</sub> represents C<sub>1-4</sub>alkyl;
- R<sub>8</sub> represents C<sub>1-4</sub>alkyl or R<sub>7</sub> and R<sub>8</sub> taken together form a 5-7 member ring;
- m is 0, 1, 2 or 3;
- n is 0, 1, 2, or 3
- 20 p is 1, 2 or 3 and
- q is 1, 2, 3 or 4.

Preferred compounds useful in the present invention include:

- 2-Amino-4H-indeno[1,2b]thiophene-3-carboxylic acid amide;
- 2-Ureido-4H-indeno[1,2b]thiophene-3-carboxylic acid amide;
- 25 2-Amino-4,5-dihydro-naphtho[2,1-b]thiophene-3-carboxylic acid amide; and
- 2-Ureido-4,5-dihydro-naphtho[2,1-b]thiophene-3-carboxylic acid amide.

This invention provides methods for treating a variety of diseases associated with NF-κB activation including inflammatory and tissue repair disorders; particularly



rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease) osteoarthritis, osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and ultraviolet radiation (UV)-induced skin damage; autoimmune diseases including systemic lupus erythematosus, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherosclerosis, restenosis, diabetes, glomerulonephritis, cancer, including Hodgkins disease, cachexia, inflammation associated with infection and certain viral infections, including acquired immune deficiency syndrome (AIDS), adult respiratory distress syndrome, and Ataxia Telangiectasia.

# DEFINITIONS

The present invention includes all hydrates, solvates, complexes and prodrugs of the compounds of this invention. Prodrugs are any covalently bonded compounds, which release the active parent, drug according to Formulas I and II *in vivo*. If a chiral center or another form of an isomeric center is present in a compound of the present invention, all forms of such isomer or isomers, including enantiomers and diastereomers, are intended to be covered herein. Inventive compounds containing a chiral center may be used as a racemic mixture, an enantiomerically enriched mixture, or the racemic mixture may be separated using well-known techniques and an individual enantiomer may be used alone. In cases in which compounds have unsaturated carbon-carbon double bonds, both the *cis* (Z) and *trans* (E) isomers are within the scope of this invention. In cases wherein compounds may exist in tautomeric forms, such as keto-enol tautomers, each tautomeric form is contemplated as being included within this invention whether existing in equilibrium or predominantly in one form.

The meaning of any substituent at any one occurrence in Formula I or any subformula thereof is independent of its meaning, or any other substituent's meaning, at any other occurrence, unless specified otherwise.

As used herein, "alkyl" refers to an optionally substituted hydrocarbon group joined by single carbon-carbon bonds and having 1-6 carbon atoms joined together.

The alkyl hydrocarbon group may be linear, branched or cyclic, saturated or

unsaturated. Substituents on optionally substituted alkyl are selected from the group consisting of aryl, OH, O-alkyl, CO, halogen, CF<sub>3</sub>, and OCF<sub>3</sub>.

As used herein, "aryl" refers to an optionally substituted aromatic group with at least one ring having a conjugated pi-electron system, containing up to two  
5 conjugated or fused ring systems. Aryl includes carbocyclic aryl, and biaryl groups, all of which may be optionally substituted. Substituents are selected from the group consisting of halogen, C<sub>1-4</sub> alkyl, NH<sub>2</sub>, OCF<sub>3</sub>, CF<sub>3</sub>, O-alkyl, S-alkyl, CN, CHO, SO<sub>2</sub>-alkyl and NO<sub>2</sub>.

As used herein, "heteroaryl" refers to an optionally substituted aromatic  
10 group with at least one ring having a conjugated pi-electron system, containing up to two conjugated or fused ring systems and 1-3 heteroatoms selected from O, S and N. Heteroaryl includes carbocyclic heteroaryl, aryl-heteroaryl and biheteroaryl groups, all of which may be optionally substituted. Preferred aryl include phenyl and naphthyl. More preferred aryl include phenyl. Preferred substituents are selected  
15 from the group consisting of halogen, C<sub>1-4</sub> alkyl, NH<sub>2</sub>, OCF<sub>3</sub>, CF<sub>3</sub>, O-alkyl, S-alkyl, CN, CHO, SO<sub>2</sub>-alkyl and NO<sub>2</sub>. Examples of heteroaryl rings included pyrrole, furan, thiophene, indole, isoindole, benzofuran, isobenzofuran, benzothiphenene, pyridine, quinoline, isoquinoline, quinolizine, pyrazole, imidazole, isoxazole, oxazole, isothiazole, thiazole, pyridazine, pyrimidine, and pyrazine.


20 As used herein "halogen" refers to include F, Cl, Br, and I.

### METHODS OF PREPARATION


#### **General Preparation:**

The general preparation of the aminothiophene analogs is shown in Schemes  
25 1-2. Morpholine is added to a stirred solution of cyanoacetamide, sulfur, and cyclic ketone in absolute ethanol. The resulting solution is stirred at room temperature or up to 60°C overnight. The solvent is then removed under *vacuo* and the residue is taken up into ethyl acetate, washed by water and brine, dried over anhydrous magnesium sulfate, filtered and concentrated under *vacuo* to give a dark brown solid.  
30 The product is then usually purified by chromatography to give the desired product.

Scheme 1



Scheme 2



5

The following Experimentals are intended to be illustrative only and not limiting in any way.

10

### Example 1

#### **Preparation of 2-Amino-4H-indeno[1,2b]thiophene-3-carboxylic acid amide trifluoroacetate**

Morpholine (1 mL) was added dropwise to a stirred solution of cyanoacetamide (0.84 g, 0.01 mol), sulfur (0.36, 0.012 mol), and 2-indanone (1.32 g, 0.01 mmol) in absolute ethanol (5 mL). The resulting solution was stirred at 60 °C overnight. The solvent was then removed under vacuo and the residue was taken up into ethyl acetate (10 mL), washed by water (2 x 10 mL) and brine (10 mL), dried over anhydrous magnesium sulfate, filtered and concentrated under vacuo to give a

dark brown solid. The product was then purified on Gilson preparative HPLC (YMC HPLC column 50 X 20 mm I.D., s-5  $\mu$ m, 120 Å; gradient elution, 0.1%TFA in acetonitrile:0.1% aqueous TFA, 10:90 to 90:10, 10 min) to give the title compound as a brown solid (100 mg, 0.435 mmol, 4.3% yield). ESMS m/z: 231

5 [M+H]<sup>+</sup>.

### Example 2

**Preparation of 2-Ureido-4H-indeno[1,2b]thiophene-3-carboxylic acid amide**  
Chlorosulfonyl isocyanate (0.025 g, 0.17 mL) was added dropwise to a

10 stirred solution of 2-amino-4H-indeno[1,2b]thiophene-3-carboxylic acid amide trifluoroacetate (0.040 g, 0.17 mmol) in dry dichloromethane (2 mL). The resulting reaction mixture was stirred under nitrogen for 30 min. Water (0.5 mL) was then added to the reaction mixture and the reaction mixture was allowed to stir an additional 10 minutes before the solvent was removed under vacuo. The residue was

15 then purified on Gilson preparative HPLC (YMC HPLC column 50 X 20 mm I.D., s-5  $\mu$ m, 120 Å; gradient elution, 0.1%TFA in acetonitrile: 0.1% aqueous TFA, 10:90 to 90:10, 10 min) to give title compound as brown solid (0.020 g, 0.435 mmol, 43.5% yield). ESMS m/z: 274 [M+H]<sup>+</sup>.

20

### Example 3

**Preparation of 2-Amino-4,5-dihydro-naphtho[2,1-d]thiophene-3-carboxylic acid amide**  
The title compound was prepared by the same procedure as Example 1

25 except that 2-indanone was replaced with beta-tetralone to give the above titled compound as a brown solid (60 % pure by LC-MS). ESMS m/z: 245 [M+H]<sup>+</sup>.

### Example 4

30 **Preparation of 2-Ureido-4,5-dihydro-naphtho[2,1-b]thiophene-3-carboxylic acid amide**  
The title compound was prepared by the same procedure as Example 2 except 2-amino-4H-indeno[1,2b]thiophene-3-carboxylic acid amide was replaced

with 2-amino-4,5-dihydro-naphtho[1,2-d]thiophene-3-carboxylic acid amide to give the above titled compound as a brown solid. ESMS  $m/z$ : 288  $[M+H]^+$ .

5 This invention provides a pharmaceutical composition, which comprises a compound according to Formula, I and a pharmaceutically acceptable carrier, diluent or excipient. Accordingly, the compounds of Formula I may be used in the manufacture of a medicament. Pharmaceutical compositions of the compounds of Formula I prepared as hereinbefore described may be formulated as solutions or lyophilized powders for parenteral administration. Powders may be reconstituted by  
10 addition of a suitable diluent or other pharmaceutically acceptable carrier prior to use. The liquid formulation may be a buffered, isotonic, aqueous solution. Examples of suitable diluents are normal isotonic saline solution, standard 5% dextrose in water or buffered sodium or ammonium acetate solution. Such formulation is especially suitable for parenteral administration, but may also be used  
15 for oral administration or contained in a metered dose inhaler or nebulizer for insufflation. It may be desirable to add excipients such as polyvinylpyrrolidone, gelatin, hydroxy cellulose, acacia, polyethylene glycol, mannitol, sodium chloride or sodium citrate.

Alternately, these compounds may be encapsulated, tableted or prepared in  
20 an emulsion or syrup for oral administration. Pharmaceutically acceptable solid or liquid carriers may be added to enhance or stabilize the composition, or to facilitate preparation of the composition. Solid carriers include starch, lactose, calcium sulfate dihydrate, terra alba, magnesium stearate or stearic acid, talc, pectin, acacia, agar or gelatin. Liquid carriers include syrup, peanut oil, olive oil, saline and water.  
25 The carrier may also include a sustained release material such as glyceryl monostearate or glyceryl distearate, alone or with a wax. The amount of solid carrier varies but, preferably, will be between about 20 mg to about 1 g per dosage unit. The pharmaceutical preparations are made following the conventional techniques of pharmacy involving milling, mixing, granulating, and compressing,  
30 when necessary, for tablet forms; or milling, mixing and filling for hard gelatin capsule forms. When a liquid carrier is used, the preparation will be in the form of a

syrup, elixir, emulsion or an aqueous or non-aqueous suspension. Such a liquid formulation may be administered directly p.o. or filled into a soft gelatin capsule.

For rectal administration, the compounds of this invention may also be combined with excipients such as cocoa butter, glycerin, gelatin or polyethylene glycols and molded into a suppository.

The methods of the present invention include topical inhaled and intracolonic administration of the compounds of Formula I. By topical administration is meant non-systemic administration, including the application of a compound of the invention externally to the epidermis, to the buccal cavity and instillation of such a compound into the ear, eye and nose, wherein the compound does not significantly enter the blood stream. By systemic administration is meant oral, intravenous, intraperitoneal and intramuscular administration. The amount of a compound of the invention (hereinafter referred to as the active ingredient) required for therapeutic or prophylactic effect upon topical administration will, of course, vary with the compound chosen, the nature and severity of the condition being treated and the animal undergoing treatment, and is ultimately at the discretion of the physician

While it is possible for an active ingredient to be administered alone as the raw chemical, it is preferable to present it as a pharmaceutical formulation. The active ingredient may comprise, for topical administration, from 0.01 to 5.0 wt% of the formulation.

The topical formulations of the present invention, both for veterinary and for human medical use, comprise an active ingredient together with one or more acceptable carriers therefor and optionally any other therapeutic ingredients. The carrier must be "acceptable" in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin to the site of where treatment is required such as: liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose.

Drops according to the present invention may comprise sterile aqueous or oily solutions or suspensions and may be prepared by dissolving the active

ingredient in a suitable aqueous solution of a bactericidal and/or fungicidal agent and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a suitable container, which is then sealed and sterilized by autoclaving, or maintaining  
5 at 90-100 C for half an hour. Alternatively, the solution may be sterilized by filtration and transferred to the container by an aseptic technique. Examples of bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.002%), benzalkonium chloride (0.01%) and chlorhexidine acetate (0.01%). Suitable solvents for the preparation of an oily  
10 solution include glycerol, diluted alcohol and propylene glycol.

Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods similar to those for the preparation of drops. Lotions or liniments for application to the skin may  
15 also include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil.

Creams, ointments or pastes according to the present invention are semi-solid formulations of the active ingredient for external application. They may be made by  
20 mixing the active ingredient in finely divided or powdered form, alone or in solution or suspension in an aqueous or non-aqueous fluid, with the aid of suitable machinery, with a greasy or non-greasy basis. The basis may comprise hydrocarbons such as hard, soft or liquid paraffin, glycerol, beeswax, a metallic soap, a mucilage, an oil of natural origin such as almond, corn, arachis, castor or  
25 olive oil, wool fat or its derivatives, or a fatty acid such as stearic or oleic acid together with an alcohol such as propylene glycol or macrogols. The formulation may incorporate any suitable surface active agent such as an anionic, cationic or non-ionic surface active agent such as sorbitan esters or polyoxyethylene derivatives thereof. Suspending agents such as natural gums, cellulose derivatives or in organic  
30 materials such as siliceous silicas, and other ingredients such as lanolin, may also be included.

### UTILITY OF THE PRESENT INVENTION

The compounds of Formula I are useful as inhibitors of the IKK-beta kinase phosphorylation of I $\kappa$ B and as such are inhibitors of NF- $\kappa$ B activation. The present method utilizes compositions and formulations of said compounds, including pharmaceutical compositions and formulations of said compounds.

The present invention particularly provides methods of treatment of diseases associated with inappropriate NF- $\kappa$ B activation, which methods comprise administering to an animal, particularly a mammal, most particularly a human in need thereof one or more compounds of Formula I. The present invention particularly provides methods for treating inflammatory and tissue repair disorders, particularly rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease) osteoarthritis, osteoporosis and fibrotic diseases; dermatosis, including psoriasis, atopic dermatitis and ultraviolet radiation (UV)-induced skin damage, autoimmune diseases including systemic lupus erythematosus, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherosclerosis, restenosis, diabetes, glomerulonephritis, cancer, including Hodgkins disease, cachexia, inflammation associated with infection and certain viral infections, including acquired immune deficiency syndrome (AIDS), adult respiratory distress syndrome and Ataxia Telangiectasia.

For acute therapy, parenteral administration of one or more compounds of Formula I is useful. An intravenous infusion of the compound in 5% dextrose in water or normal saline, or a similar formulation with suitable excipients, is most effective, although an intramuscular bolus injection is also useful. Typically, the parenteral dose will be about 0.01 to about 50 mg/kg; preferably between 0.1 and 20 mg/kg, in a manner to maintain the concentration of drug in the plasma at a concentration effective to inhibit IKK-beta and therefore activation of NF- $\kappa$ B. The compounds are administered one to four times daily at a level to achieve a total daily dose of about 0.4 to about 80 mg/kg/day. The precise amount of a compound used in the present method which is therapeutically effective, and the route by which such compound is



best administered, is readily determined by one of ordinary skill in the art by comparing the blood level of the agent to the concentration required to have a therapeutic effect.

The compounds of Formulas I may also be administered orally to the patient, in a manner such that the concentration of drug is sufficient to inhibit IKK-beta and therefore activation of NF- $\kappa$ B or to achieve any other therapeutic indication as disclosed herein. Typically, a pharmaceutical composition containing the compound is administered at an oral dose of between about 0.1 to about 50 mg/kg in a manner consistent with the condition of the patient. Preferably the oral dose would be about 0.5 to about 20 mg/kg.

The compounds of Formulas I may also be administered topically to the patient, in a manner such that the concentration of drug is sufficient to inhibit IKK-beta and therefore activation of NF- $\kappa$ B or to achieve any other therapeutic indication as disclosed herein. Typically, a pharmaceutical composition containing the compound is administered in a topical formulation of between about 0.01% to about 5% w/w.

No unacceptable toxicological effects are expected when compounds of the present invention are administered in accordance with the present invention.

The ability of the compounds described herein to inhibit the activation of NF- $\kappa$ B is clearly evidenced in their ability to inhibit the phosphorylation of the N-terminal fragment of I $\kappa$ B- $\alpha$  by IKK- $\beta$  (see Table 1 for examples). These compounds also block the degradation of I $\kappa$ B- $\alpha$  and the nuclear translocation of NF- $\kappa$ B in human monocytes and other mammalian cells upon activation of the cells with a pro-inflammatory stimuli (e.g., TNF- $\alpha$ , LPS, etc.). In addition these compounds inhibit pro-inflammatory mediator production from LPS-stimulated human monocytes and stimulated human primary synovial fibroblasts. The utility of the present NF- $\kappa$ B inhibitors in the therapy of diseases is premised on the importance of NF- $\kappa$ B activation in a variety of diseases.

NF- $\kappa$ B plays a key role in the regulated expression of a large number of pro-inflammatory mediators including cytokines such as TNF, IL-1 $\beta$ , IL-6 and IL-8 (Mukaida *et al.*, 1990; Liberman and Baltimore, 1990; Matsusaka *et al.*, 1993), cell

adhesion molecules, such as ICAM and VCAM (Marui *et al.*, 1993; Kawai *et al.*, 1995; Ledebur and Parks, 1995), and inducible nitric oxide synthase (iNOS) (Xie *et al.*, 1994; Adcock *et al.*, 1994). (Full reference citations are at the end of this section). Such mediators are known to play a role in the recruitment of leukocytes at sites of inflammation and in the case of iNOS, may lead to organ destruction in some inflammatory and autoimmune diseases (McCartney-Francis *et al.*, 1993; Kleemann *et al.*, 1993).

Evidence for an important role of NF- $\kappa$ B in inflammatory disorders is obtained in studies of asthmatic patients. Bronchial biopsies taken from mild atopic asthmatics show significant increases in the number of cells in the submucosa staining for activated NF- $\kappa$ B, total NF- $\kappa$ B, and NF- $\kappa$ B-regulated cytokines such as GM-CSF and TNF $\alpha$  compared to biopsies from normal non-atopic controls (Wilson *et al.*, 1998). Furthermore, the percentage of vessels expressing NF- $\kappa$ B immunoreactivity is increased as is IL-8 immunoreactivity in the epithelium of the biopsy specimens (Wilson *et al.*, 1998). As such, inhibition of IL-8 production through the inhibition of NF- $\kappa$ B, as has been demonstrated by these compounds would be predicted be beneficial in airway inflammation.

Recent studies suggest that NF- $\kappa$ B may also play a critical role in the pathogenesis of inflammatory bowel disease (IBD). Activated NF- $\kappa$ B is seen in colonic biopsy specimens from Chron's disease and ulcerative colitis patients (Ardite *et al.*, 1998; Rogler *et al.*, 1998; Schreiber *et al.*, 1998). Activation is evident in the inflamed mucosa but not in uninfamed mucosa (Ardite *et al.*, 1998; Rogler *et al.*, 1998) and is associated with increased IL-8 mRNA expression in the same sites (Ardite *et al.*, 1998). Furthermore, corticosteroid treatment strongly inhibits intestinal NF- $\kappa$ B activation and reduces colonic inflammation (Ardite *et al.*, 1998; Schreiber *et al.*, 1998). Again, inhibition of IL-8 production through the inhibition of NF- $\kappa$ B, as has been demonstrated by these compounds would be predicted be beneficial in inflammatory bowel disease.

Animal models of gastrointestinal inflammation provide further support for NF- $\kappa$ B as a key regulator of colonic inflammation. Increased NF- $\kappa$ B activity is

observed in the lamina propria macrophages in 2,4,6,-trinitrobenzene sulfonic acid (TNBS)-induced colitis in mice with p65 being a major component of the activated complexes (Neurath *et al.*, 1996; Neurath and Pettersson, 1997). Local administration of p65 antisense abrogates the signs of established colitis in the treated animals with no signs of toxicity (Neurath *et al.*, 1996; Neurath and Pettersson, 1997). As such, one would predict that small molecule inhibitors of NF- $\kappa$ B would be useful in the treatment of IBD.

Further evidence for a role of NF- $\kappa$ B in inflammatory disorders comes from studies of rheumatoid synovium. Although NF- $\kappa$ B is normally present as an inactive cytoplasmic complex, recent immunohistochemical studies have indicated that NF- $\kappa$ B is present in the nuclei, and hence active, in the cells comprising human rheumatoid synovium (Handel *et al.*, 1995; Marok *et al.*, 1996; Sioud *et al.*, 1998) and in animal models of the disease (Tsao *et al.*, 1997). The staining is associated with type A synoviocytes and vascular endothelium (Marok *et al.*, 1996). Furthermore, constitutive activation of NF- $\kappa$ B is seen in cultured synoviocytes (Roshak *et al.*, 1996; Miyazawa *et al.*, 1998) and in synovial cell cultures stimulated with IL-1 $\beta$  or TNF $\alpha$  (Roshak *et al.*, 1996; Fujisawa *et al.*, 1996; Roshak *et al.*, 1997). Thus, the activation of NF- $\kappa$ B may underlie the increased cytokine production and leukocyte infiltration characteristic of inflamed synovium. The ability of these compounds to inhibit NF- $\kappa$ B and thereby inhibit the production of pro-inflammatory mediators (e.g. cytokines and prostanoids) by these cells would be predicted to yield benefit in rheumatoid arthritis.

#### **Biological Assays:**

The compounds of this invention may be tested in one of several biological assays to determine the concentration of compound, which is required to have a given pharmacological effect.

NF- $\kappa$ B activity may also be measured in an electrophoretic mobility shift assay (EMSA) to assess the presence of NF- $\kappa$ B protein in the nucleus. The cells of interest are cultured to a density of  $1 \times 10^6$  /mL. The cells are harvested by centrifugation, washed in PBS without Ca<sup>2+</sup> and Mg<sup>2+</sup> and resuspended in PBS with

- Ca<sup>2+</sup> and Mg<sup>2+</sup> at 1x10<sup>7</sup> cells/mL. To examine the effect of compound on the activation of NF-κB, the cell suspensions are treated with various concentrations of drug or vehicle (DMSO, 0.1%) for 30 min. at 37 °C prior to stimulation with TNF-α (5.0 ng/mL) for an additional 15 min. Cellular and nuclear extracts are prepared
- 5 follows. Briefly, at the end of the incubation period the cells (1x10<sup>7</sup> cells) are washed 2x in PBS without Ca<sup>2+</sup> and Mg<sup>2+</sup>. The resulting cell pellets are resuspended in 20 uL of Buffer A (10 mM Hepes (pH 7.9), 10 mM KCl, 1.5 mM MgCl<sub>2</sub>, 0.5 mM dithiothreitol (DTT) and 0.1% NP-40) and incubated on ice for 10 min. The nuclei are pelleted by microcentrifugation at 3500 rpm for 10 min at 4 °C. The resulting
- 10 supernatant was collected as the cellular extract and the nuclear pellet was resuspended in 15 uL Buffer C (20 mM Hepes (pH 7.9), 0.42 M NaCl, 1.5mM MgCl<sub>2</sub>, 25% glycerol, 0.2 mM EDTA, 0.5 mM DTT, and 0.5 mM phenylmethylsulphonyl fluoride (PMSF)). The suspensions are mixed gently for 20 min at 4 °C then microcentrifuged at 14,000 rpm for 10 min at 4 °C. The
- 15 supernatant is collected and diluted to 60 uL with Buffer D (20mM Hepes (pH 7.9), 50 mM KCl, 20% glycerol, 0.2 mM EDTA, 0.5 mM DTT, and 0.5 mM PMSF). All samples are stored at -80 °C until analyzed. The protein concentration of the extracts is determined according to the method of Bradford (Bradford, 1976) with BioRad reagents.
- 20 The effect of compounds on transcription factor activation is assessed in an electrophoretic mobility shift assay (EMSA) using nuclear extracts from treated cells as described above. The double stranded NF-κB consensus oligonucleotides (5'-AGTTGAGGGGACTTTCCCAGGC-3') are labelled with T<sub>4</sub> polynucleotide kinase and [γ-<sup>32</sup>P]ATP. The binding mixture (25 uL) contains 10 mM Hepes-NaOH (pH
- 25 7.9), 4 mM Tris-HCl (pH 7.9), 60 mM KCl, 1 mM EDTA, 1 mM dithiothreitol, 10% glycerol, 0.3 mg/mL bovine serum albumin, and 1 ug poly(dI-dC)•poly(dI-dC). The binding mixtures (10 ug nuclear extract protein) are incubated for 20 min at room temperature with 0.5 ng of <sup>32</sup>P-labelled oligonucleotide (50,000-100,000 cpm) in the presence or absence of unlabeled competitor after which the mixture is loaded on a
- 30 4% polyacrylamide gel prepared in 1X Tris borate/EDTA and electrophoresed at 200

V for 2 h. Following electrophoresis the gels are dried and exposed to film for detection of the binding reaction.

The effect of compounds on the phosphorylation of I $\kappa$ B may be monitored in a Western blot. Cellular extracts are subjected to sodium dodecyl sulfate-

- 5 polyacrylamide gel electrophoresis (SDS-PAGE) on 10% gels (BioRad, Hercules, CA) and the proteins transferred to nitrocellulose sheets (Hybond<sup>tm</sup>-ECL, Amersham Corp., Arlington Heights, IL). Immunoblot assays are performed using a polyclonal rabbit antibody directed against I $\kappa$ B $\alpha$  or I $\kappa$ B $\beta$  followed with a peroxidase-conjugated donkey anti-rabbit secondary antibody (Amersham Corp., Arlington Heights, IL). Immunoreactive bands are detected using the Enhanced Chemiluminescence (ECL) assay system (Amersham Corp., Arlington Heights, IL).

- Assays for I $\kappa$ B kinases were conducted as follows: IKK- $\alpha$  was expressed as a hexa-histidine tagged protein in baculovirus-infected insect cells and purified over a Ni-NTA affinity column. Kinase activity was assayed using 50 ng of purified  
15 protein in assay buffer (20 mM Hepes, pH 7.7, 2 mM MgCl<sub>2</sub>, 1 mM MnCl<sub>2</sub>, 10 mM  $\beta$ -glycerophosphate, 10 mM NaF, 10 mM PNPP, 0.3 mM Na<sub>3</sub>VO<sub>4</sub>, 1 mM benzamidine, 2  $\mu$ M PMSF, 10  $\mu$ g/ml aprotinin, 1  $\mu$ g/mL leupeptin, 1  $\mu$ g/mL pepstatin, 1mM DTT) containing various concentrations of compound or DMSO vehicle and ATP as indicated (Pharmacia Biotech Inc., Piscataway, NJ). The  
20 reaction was started by the addition of 200 ng I $\kappa$ B-GST (Santa Cruz Biotechnology, Inc., Santa Cruz, CA), in a total volume of 50  $\mu$ L. The reaction was allowed to proceed for 1 h. at 30 °C after which the reaction was terminated by the addition of EDTA to a final concentration of 20 mM. Kinase activity was determined by dissociation-enhanced lanthanide fluorescence immunoassay (Wallac Oy, Turku,  
25 Finland) using a phospho-I $\kappa$ B- $\alpha$  (Ser32) antibody (New England Biolabs, Inc., Beverly, MA) and an Eu<sup>3+</sup>-labelled anti-rabbit IgG (Wallac Oy, Turku, Finland). The plates were read in a VICTOR 1420 Multilabel Counter (Wallac), using a standard europium protocol (excitation 340 nm, emission 615 nm; fluorescence measured for 400  $\mu$ s after a 400  $\mu$ sec delay). Data are expressed as fluorescence  
30 (cps) units.

IKK- $\beta$  was expressed as a GST-tagged protein, and its activity was assessed in a 96-well scintillation proximity assay (SPA). Briefly, IKK- $\beta$  was diluted in assay buffer as described above (20 nM final), with various concentrations of compound or DMSO vehicle, 240 nM ATP and 200 nCi [ $\gamma$ - $^{33}$ P]-ATP (10 mCi/mL, 2000 Ci/mmol; NEN Life Science Products, Boston, MA). The reaction was started with the addition of a biotinylated peptide comprising amino acids 15 – 46 of I $\kappa$ B- $\alpha$  (American Peptide) to a final concentration of 2.4  $\mu$ M, in a total volume of 50  $\mu$ L. The sample incubated for one hour at 30  $^{\circ}$ C, followed by the addition of 150  $\mu$ L of stop buffer (PBS w/o Ca $^{2+}$ , Mg $^{2+}$ , 0.1% Triton X-100 (v/v), 10 mM EDTA) containing 0.2 mg streptavidin-coated SPA PVT beads (Amersham Pharmacia Biotech, Piscataway, NJ). The sample was mixed, incubated for 10 min. at room temperature, centrifuged (1000 xg, 2 minutes), and measured on a Hewlett-Packard TopCount.

The effect of IKK- $\beta$  inhibitors on primary synovial fibroblast mediator production was assessed as follows: Primary cultures of human RSF were obtained by enzymatic digestion of synovium obtained from adult patients with rheumatoid arthritis as previously described (Roshak *et al.*, 1996b). Cells were cultured in Earl's Minimal Essential Medium (EMEM) which contained 10% fetal bovine serum (FBS), 100 units/ml penicillin and 100  $\mu$ g/ml streptomycin (GIBCO, Grand Island, NY), at 37 $^{\circ}$ C and 5% CO $_2$ . Cultures were used at passages 4 through 9 in order to obtain a more uniform type B fibroblast population. For some studies, fibroblasts were plated at 5 x 10 $^4$  cells/mL in 16 mm (diameter) 24 well plates (Costar, Cambridge, MA). Cells (70-80% confluence) were exposed to IL-1 $\beta$  (1 ng/mL) (Genzyme, Cambridge, MA) for the designated time. Drugs in DMSO vehicle (1%) were added to the cell cultures 15 minutes prior to the addition of IL-1. Studies were conducted 3-4 times using synovial cells from different donors. RSF cellular extracts were prepared from cells treated as described above. Briefly, human RSF were removed by trypsin/EDTA, washed, and harvested by centrifugation. Cellular extracts were prepared as previously described (Dignam *et al.*, 1983; Osborn, *et al.*, 1989). Briefly, at the end of the incubation period the cells (1x10 $^7$  cells) were washed 2x in PBS without Ca $^{2+}$  and Mg $^{2+}$ . The resulting cell pellets were

resuspended in 20  $\mu$ L of Buffer A (10 mM Hepes (pH 7.9), 10 mM KCl, 1.5 mM  $\text{MgCl}_2$ , 0.5 mM.

Effect of IKK- $\beta$  inhibition on human monocyte stimulated eicosanoid and cytokine production was assessed as follows: Monocytes were isolated from  
5 heparinized whole blood by double gradient centrifugation as previously described. Isolated monocyte enriched PBMCs were then adhered to 24 well culture plates at  $2 \times 10^6$  cells/mL in RPMI 1640 10% FBS (Hyclone, Logan, Utah) for 2 h. to further enrich the monocyte population. The media was then removed, cells washed once with RPMI 1640, and 1 mL RPMI 1640 10% FBS was added to the wells. Test  
10 compounds are added to the wells with a final vehicle concentration of 0.05% DMSO. Monocytes were activated by the addition of 200 ng/mL endotoxin (LPS; *E. coli* serotype 026:B6)(Sigma, St. Louis, MO.) and incubated for 24 hrs. Cell-free supernates were analyzed by ELISA for TNF- $\alpha$  (EIA developed at SB), PGE<sub>2</sub> (Cayman Chemical, Ann Arbor, MI), and IL-8 and IL-6 Biosource International,  
15 Camarillo, CA). Viability of the cells was determined by trypan blue exclusion.

Effect of IKK- $\beta$  inhibitors on phorbol ester-induced inflammation was assessed as follows: The inflammatory response induced by the cutaneous application of phorbol ester (PMA) to the external pinnae of Balb/c mice has proven to be a useful model to examine multifactorial inflammatory cell infiltration and  
20 inflammatory alteration of epidermis. The intense inflammatory lesion is dominated by neutrophil infiltration, which can be easily quantified by measurement tissue concentration myeloperoxidase, an azuriphilic granular enzyme present in neutrophils. In addition, the overall intensity of the inflammatory response can be measured by determination of ear thickness. Balb/c mice (n = 6/group) were  
25 administered drug treatment or vehicle followed by PMA (4  $\mu$ g/ear). The mice were sacrificed 4 h. later, the ear thickness determined and NF- $\kappa$ B activation was monitored by IkB $\alpha$  western or EMSA analysis.

Effect of IKK- $\beta$  inhibitors on rat carrageenan-induced paw edema was assessed as follows: Male Lewis rats (Charles River- Raleigh, NC) were housed and  
30 allowed free access to food and water, and weighed between 200-275g for each experiment. Compound or vehicle (0.5% Tragacanth (p.o.) or 10%DMSO,

- 5%DMA, 30% Cremophor(i.p.)) was administered 30 minutes to 1 hour prior to the carrageenan injection. Edema was induced by injection of 1% carrageenan in sterile dH<sub>2</sub>O (0.05ml/paw) into the plantar surface of the right hindpaw. Paw thickness was measured prior to administration of compound or vehicle, and again at 3 hours, to determine change in paw volume. Rats were euthanized by CO<sub>2</sub> inhalation and the right hindfoot was removed, immediately frozen in liquid nitrogen and stored at -80C for analysis.

### General

- Nuclear magnetic resonance spectra were recorded at either 250, 300 or 400 MHz using, respectively, a Bruker AM 250, Bruker ARX 300 or Bruker AC 400 spectrometer. CDCl<sub>3</sub> is deuteriochloroform, DMSO-d<sub>6</sub> is hexadeuteriodimethylsulfoxide, and CD<sub>3</sub>OD is tetradeuteriomethanol. Chemical shifts are reported in parts per million (δ) downfield from the internal standard tetramethylsilane. Abbreviations for NMR data are as follows: s = singlet, d = doublet, t = triplet, q = quartet, m = multiplet, dd = doublet of doublets, dt = doublet of triplets, app = apparent, br = broad. J indicates the NMR coupling constant measured in Hertz. Continuous wave infrared (IR) spectra were recorded on a Perkin-Elmer 683 infrared spectrometer, and Fourier transform infrared (FTIR) spectra were recorded on a Nicolet Impact 400 D infrared spectrometer. IR and FTIR spectra were recorded in transmission mode, and band positions are reported in inverse wavenumbers (cm<sup>-1</sup>). Mass spectra were taken on either VG 70 FE, PE Syx API III, or VG ZAB HF instruments, using fast atom bombardment (FAB) or electrospray (ES) ionization techniques. Elemental analyses were obtained using a Perkin-Elmer 240C elemental analyzer. Melting points were taken on a Thomas-Hoover melting point apparatus and are uncorrected. All temperatures are reported in degrees Celsius.

Analtech Silica Gel GF and E. Merck Silica Gel 60 F-254 thin layer plates were used for thin layer chromatography. Both flash and gravity chromatography were carried out on E. Merck Kieselgel 60 (230-400 mesh) silica gel.

Where indicated, certain of the materials were purchased from the Aldrich Chemical Co., Milwaukee, Wisconsin, TCI America, Portland, OR..






- 5

What is claimed is:

1. A compound according to formula (I) hereinbelow:



(I)

5

wherein:

R<sub>1</sub> represents NR<sub>4</sub>R<sub>5</sub>;

R<sub>2</sub> represents CONH<sub>2</sub> or SO<sub>2</sub>NH<sub>2</sub>;

- 10 R<sub>3</sub> is selected from the group consisting of halogen, C<sub>1-4</sub>alkyl, NH<sub>2</sub>, CF<sub>3</sub>, OCF<sub>3</sub>, O-alkyl, S-alkyl, CN, CHO, SO<sub>2</sub>-alkyl, and NO<sub>2</sub>;

R<sub>4</sub> represents C<sub>1-4</sub>alkyl;

R<sub>5</sub> represents CONHR<sub>6</sub>;

R<sub>6</sub> is selected from the group consisting of hydrogen, alkyl and aryl;

m is 0 or 1;

- 15 n is 1 or 2; and

p is 1, 2 or 3.

2. A compound according to claim 1 wherein the compound is selected from the group consisting of:

- 2-Amino-4H-indeno[1,2b]thiophene-3-carboxylic acid amide;  
 20 2-Ureido-4H-indeno[1,2b]thiophene-3-carboxylic acid amide;  
 2-Amino-4,5-dihydro-naphtho[2,1-b]thiophene-3-carboxylic acid amide; and  
 2-Ureido-4,5-dihydro-naphtho[2,1-b]thiophene-3-carboxylic acid amide.

3. A method of treating a disease characterized by pathological NF-κB  
 25 activation comprising inhibiting the pathological activation by administering to a patient in need thereof an effective amount of a compound according to claim 1.

5  
10  
15  
20  
25  
30

4. A method according to claim 3 wherein the disease is an inflammatory or tissue repair disorder.

5. A method according to Claim 4 wherein the disease is selected from the group consisting of inflammatory and tissue repair disorders, particularly rheumatoid arthritis, inflammatory bowel disease, asthma and COPD (chronic obstructive pulmonary disease) osteoarthritis, osteoporosis and fibrotic diseases, dermatosis, including psoriasis, atopic dermatitis and ultraviolet radiation (UV)-induced skin damage, autoimmune diseases including systemic lupus erythematosus, multiple sclerosis, psoriatic arthritis, ankylosing spondylitis, tissue and organ rejection, Alzheimer's disease, stroke, atherosclerosis, restenosis, diabetes, glomerulonephritis, cancer, including Hodgkins disease, cachexia, inflammation associated with infection and certain viral infections, including acquired immune deficiency syndrome (AIDS), adult respiratory distress syndrome, and Ataxia Telangiectasia.

6. A method according to Claim 3 wherein said disease is dermatosis.

7. A method according to Claim 3 wherein the disease is selected from the group consisting of: psoriasis, atopic dermatitis, and UV-induced skin damage.

8. A method according to Claim 3 wherein the disease is selected from the group consisting of autoimmune diseases; tissue and organ rejection, Alzheimer's disease, stroke, atherosclerosis, restenosis, diabetes, glomerulonephritis, osteoarthritis, osteoporosis, and Ataxia Telangiectasia.

9. A method according to Claim 3 wherein said disease is an autoimmune disease.

10. A method according to Claim 3 wherein the autoimmune disease is systemic lupus erythematosus, multiple sclerosis, psoriatic arthritis, or ankylosing spondylitis, diabetes

11. A method according to any one of Claim 1 wherein the disease is cancer and or cachexia.
- 5 12. A method according to Claim 3 wherein the cancer is Hodgkins disease.
13. A method according to Claim 3 wherein the disease is inflammation associated with infection and certain viral infections, including acquired immune deficiency syndrome (AIDS).
- 10 14. A method according to Claim 3 wherein the disease is AIDS.
15. A method according to Claim 3 wherein the disease is adult respiratory distress syndrome.
- 15 16. A method according to claim 3 wherein there is dual inhibition of NF- $\kappa$ B and checkpoint kinase.

# ABSTRACT OF THE INVENTION

The present invention provides novel compounds and methods for using  
5 them to treat diseases with aminothiophene inhibitors of IKK- $\beta$  phosphorylation of  
I $\kappa$ B. In so doing these aminothiophene inhibitors block pathological activation of  
transcription factor NF- $\kappa$ B in which diseases excessive activation of NF- $\kappa$ B is  
implicated.

20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100 101 102 103 104 105 106 107 108 109 110 111 112 113 114 115 116 117 118 119 120 121 122 123 124 125 126 127 128 129 130 131 132 133 134 135 136 137 138 139 140 141 142 143 144 145 146 147 148 149 150 151 152 153 154 155 156 157 158 159 160 161 162 163 164 165 166 167 168 169 170 171 172 173 174 175 176 177 178 179 180 181 182 183 184 185 186 187 188 189 190 191 192 193 194 195 196 197 198 199 200 201 202 203 204 205 206 207 208 209 210 211 212 213 214 215 216 217 218 219 220 221 222 223 224 225 226 227 228 229 230 231 232 233 234 235 236 237 238 239 240 241 242 243 244 245 246 247 248 249 250 251 252 253 254 255 256 257 258 259 260 261 262 263 264 265 266 267 268 269 270 271 272 273 274 275 276 277 278 279 280 281 282 283 284 285 286 287 288 289 290 291 292 293 294 295 296 297 298 299 300 301 302 303 304 305 306 307 308 309 310 311 312 313 314 315 316 317 318 319 320 321 322 323 324 325 326 327 328 329 330 331 332 333 334 335 336 337 338 339 340 341 342 343 344 345 346 347 348 349 350 351 352 353 354 355 356 357 358 359 360 361 362 363 364 365 366 367 368 369 370 371 372 373 374 375 376 377 378 379 380 381 382 383 384 385 386 387 388 389 390 391 392 393 394 395 396 397 398 399 400 401 402 403 404 405 406 407 408 409 410 411 412 413 414 415 416 417 418 419 420 421 422 423 424 425 426 427 428 429 430 431 432 433 434 435 436 437 438 439 440 441 442 443 444 445 446 447 448 449 450 451 452 453 454 455 456 457 458 459 460 461 462 463 464 465 466 467 468 469 470 471 472 473 474 475 476 477 478 479 480 481 482 483 484 485 486 487 488 489 490 491 492 493 494 495 496 497 498 499 500 501 502 503 504 505 506 507 508 509 510 511 512 513 514 515 516 517 518 519 520 521 522 523 524 525 526 527 528 529 530 531 532 533 534 535 536 537 538 539 540 541 542 543 544 545 546 547 548 549 550 551 552 553 554 555 556 557 558 559 560 561 562 563 564 565 566 567 568 569 570 571 572 573 574 575 576 577 578 579 580 581 582 583 584 585 586 587 588 589 590 591 592 593 594 595 596 597 598 599 600 601 602 603 604 605 606 607 608 609 610 611 612 613 614 615 616 617 618 619 620 621 622 623 624 625 626 627 628 629 630 631 632 633 634 635 636 637 638 639 640 641 642 643 644 645 646 647 648 649 650 651 652 653 654 655 656 657 658 659 660 661 662 663 664 665 666 667 668 669 670 671 672 673 674 675 676 677 678 679 680 681 682 683 684 685 686 687 688 689 690 691 692 693 694 695 696 697 698 699 700 701 702 703 704 705 706 707 708 709 710 711 712 713 714 715 716 717 718 719 720 721 722 723 724 725 726 727 728 729 730 731 732 733 734 735 736 737 738 739 740 741 742 743 744 745 746 747 748 749 750 751 752 753 754 755 756 757 758 759 760 761 762 763 764 765 766 767 768 769 770 771 772 773 774 775 776 777 778 779 780 781 782 783 784 785 786 787 788 789 790 791 792 793 794 795 796 797 798 799 800 801 802 803 804 805 806 807 808 809 810 811 812 813 814 815 816 817 818 819 820 821 822 823 824 825 826 827 828 829 830 831 832 833 834 835 836 837 838 839 840 841 842 843 844 845 846 847 848 849 850 851 852 853 854 855 856 857 858 859 860 861 862 863 864 865 866 867 868 869 870 871 872 873 874 875 876 877 878 879 880 881 882 883 884 885 886 887 888 889 890 891 892 893 894 895 896 897 898 899 900 901 902 903 904 905 906 907 908 909 910 911 912 913 914 915 916 917 918 919 920 921 922 923 924 925 926 927 928 929 930 931 932 933 934 935 936 937 938 939 940 941 942 943 944 945 946 947 948 949 950 951 952 953 954 955 956 957 958 959 960 961 962 963 964 965 966 967 968 969 970 971 972 973 974 975 976 977 978 979 980 981 982 983 984 985 986 987 988 989 990 991 992 993 994 995 996 997 998 999 1000 1001 1002 1003 1004 1005 1006 1007 1008 1009 1010 1011 1012 1013 1014 1015 1016 1017 1018 1019 1020 1021 1022 1023 1024 1025 1026 1027 1028 1029 1030 1031 1032 1033 1034 1035 1036 1037 1038 1039 1040 1041 1042 1043 1044 1045 1046 1047 1048 1049 1050 1051 1052 1053 1054 1055 1056 1057 1058 1059 1060 1061 1062 1063 1064 1065 1066 1067 1068 1069 1070 1071 1072 1073 1074 1075 1076 1077 1078 1079 1080 1081 1082 1083 1084 1085 1086 1087 1088 1089 1090 1091 1092 1093 1094 1095 1096 1097 1098 1099 1100 1101 1102 1103 1104 1105 1106 1107 1108 1109 1110 1111 1112 1113 1114 1115 1116 1117 1118 1119 1120 1121 1122 1123 1124 1125 1126 1127 1128 1129 1130 1131 1132 1133 1134 1135 1136 1137 1138 1139 1140 1141 1142 1143 1144 1145 1146 1147 1148 1149 1150 1151 1152 1153 1154 1155 1156 1157 1158 1159 1160 1161 1162 1163 1164 1165 1166 1167 1168 1169 1170 1171 1172 1173 1174 1175 1176 1177 1178 1179 1180 1181 1182 1183 1184 1185 1186 1187 1188 1189 1190 1191 1192 1193 1194 1195 1196 1197 1198 1199 1200 1201 1202 1203 1204 1205 1206 1207 1208 1209 1210 1211 1212 1213 1214 1215 1216 1217 1218 1219 1220 1221 1222 1223 1224 1225 1226 1227 1228 1229 1230 1231 1232 1233 1234 1235 1236 1237 1238 1239 1240 1241 1242 1243 1244 1245 1246 1247 1248 1249 1250 1251 1252 1253 1254 1255 1256 1257 1258 1259 1260 1261 1262 1263 1264 1265 1266 1267 1268 1269 1270 1271 1272 1273 1274 1275 1276 1277 1278 1279 1280 1281 1282 1283 1284 1285 1286 1287 1288 1289 1290 1291 1292 1293 1294 1295 1296 1297 1298 1299 1300 1301 1302 1303 1304 1305 1306 1307 1308 1309 1310 1311 1312 1313 1314 1315 1316 1317 1318 1319 1320 1321 1322 1323 1324 1325 1326 1327 1328 1329 1330 1331 1332 1333 1334 1335 1336 1337 1338 1339 1340 1341 1342 1343 1344 1345 1346 1347 1348 1349 1350 1351 1352 1353 1354 1355 1356 1357 1358 1359 1360 1361 1362 1363 1364 1365 1366 1367 1368 1369 1370 1371 1372 1373 1374 1375 1376 1377 1378 1379 1380 1381 1382 1383 1384 1385 1386 1387 1388 1389 1390 1391 1392 1393 1394 1395 1396 1397 1398 1399 1400 1401 1402 1403 1404 1405 1406 1407 1408 1409 1410 1411 1412 1413 1414 1415 1416 1417 1418 1419 1420 1421 1422 1423 1424 1425 1426 1427 1428 1429 1430 1431 1432 1433 1434 1435 1436 1437 1438 1439 1440 1441 1442 1443 1444 1445 1446 1447 1448 1449 1450 1451 1452 1453 1454 1455 1456 1457 1458 1459 1460 1461 1462 1463 1464 1465 1466 1467 1468 1469 1470 1471 1472 1473 1474 1475 1476 1477 1478 1479 1480 1481 1482 1483 1484 1485 1486 1487 1488 1489 1490 1491 1492 1493 1494 1495 1496 1497 1498 1499 1500 1501 1502 1503 1504 1505 1506 1507 1508 1509 1510 1511 1512 1513 1514 1515 1516 1517 1518 1519 1520 1521 1522 1523 1524 1525 1526 1527 1528 1529 1530 1531 1532 1533 1534 1535 1536 1537 1538 1539 1540 1541 1542 1543 1544 1545 1546 1547 1548 1549 1550 1551 1552 1553 1554 1555 1556 1557 1558 1559 1560 1561 1562 1563 1564 1565 1566 1567 1568 1569 1570 1571 1572 1573 1574 1575 1576 1577 1578 1579 1580 1581 1582 1583 1584 1585 1586 1587 1588 1589 1590 1591 1592 1593 1594 1595 1596 1597 1598 1599 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619 1620 1621 1622 1623 1624 1625 1626 1627 1628 1629 1630 1631 1632 1633 1634 1635 1636 1637 1638 1639 1640 1641 1642 1643 1644 1645 1646 1647 1648 1649 1650 1651 1652 1653 1654 1655 1656 1657 1658 1659 1660 1661 1662 1663 1664 1665 1666 1667 1668 1669 1670 1671 1672 1673 1674 1675 1676 1677 1678 1679 1680 1681 1682 1683 1684 1685 1686 1687 1688 1689 1690 1691 1692 1693 1694 1695 1696 1697 1698 1699 1700 1701 1702 1703 1704 1705 1706 1707 1708 1709 1710 1711 1712 1713 1714 1715 1716 1717 1718 1719 1720 1721 1722 1723 1724 1725 1726 1727 1728 1729 1730 1731 1732 1733 1734 1735 1736 1737 1738 1739 1740 1741 1742 1743 1744 1745 1746 1747 1748 1749 1750 1751 1752 1753 1754 1755 1756 1757 1758 1759 1760 1761 1762 1763 1764 1765 1766 1767 1768 1769 1770 1771 1772 1773 1774 1775 1776 1777 1778 1779 1780 1781 1782 1783 1784 1785 1786 1787 1788 1789 1790 1791 1792 1793 1794 1795 1796 1797 1798 1799 1800 1801 1802 1803 1804 1805 1806 1807 1808 1809 1810 1811 1812 1813 1814 1815 1816 1817 1818 1819 1820 1821 1822 1823 1824 1825 1826 1827 1828 1829 1830 1831 1832 1833 1834 1835 1836 1837 1838 1839 1840 1841 1842 1843 1844 1845 1846 1847 1848 1849 1850 1851 1852 1853 1854 1855 1856 1857 1858 1859 1860 1861 1862 1863 1864 1865 1866 1867 1868 1869 1870 1871 1872 1873 1874 1875 1876 1877 1878 1879 1880 1881 1882 1883 1884 1885 1886 1887 1888 1889 1890 1891 1892 1893 1894 1895 1896 1897 1898 1899 1900 1901 1902 1903 1904 1905 1906 1907 1908 1909 1910 1911 1912 1913 1914 1915 1916 1917 1918 1919 1920 1921 1922 1923 1924 1925 1926 1927 1928 1929 1930 1931 1932 1933 1934 1935 1936 1937 1938 1939 1940 1941 1942 1943 1944 1945 1946 1947 1948 1949 1950 1951 1952 1953 1954 1955 1956 1957 1958 1959 1960 1961 1962 1963 1964 1965 1966 1967 1968 1969 1970 1971 1972 1973 1974 1975 1976 1977 1978 1979 1980 1981 1982 1983 1984 1985 1986 1987 1988 1989 1990 1991 1992 1993 1994 1995 1996 1997 1998 1999 2000 2001 2002 2003 2004 2005 2006 2007 2008 2009 2010 2011 2012 2013 2014 2015 2016 2017 2018 2019 2020 2021 2022 2023 2024 2025 2026 2027 2028 2029 2030 2031 2032 2033 2034 2035 2036 2037 2038 2039 2040 2041 2042 2043 2044 2045 2046 2047 2048 2049 2050 2051 2052 2053 2054 2055 2056 2057 2058 2059 2060 2061 2062 2063 2064 2065 2066 2067 2068 2069 2070 2071 2072 2073 2074 2075 2076 2077 2078 2079 2080 2081 2082 2083 2084 2085 2086 2087 2088 2089 2090 2091 2092 2093 2094 2095 2096 2097 2098 2099 2100 2101 2102 2103 2104 2105 2106 2107 2108 2109 2110 2111 2112 2113 2114 2115 2116 2117 2118 2119 2120 2121 2122 2123 2124 2125 2126 2127 2128 2129 2130 2131 2132 2133 2134 2135 2136 2137 2138 2139 2140 2141 2142 2143 2144 2145 2146 2147 2148 2149 2150 2151 2152 2153 2154 2155 2156 2157 2158 2159 2160 2161 2162 2163 2164 2165 2166 2167 2168 2169 2170 2171 2172 2173 2174 2175 2176 2177 2178 2179 2180 2181 2182 2183 2184 2185 2186 2187 2188 2189 2190 2191 2192 2193 2194 2195 2196 2197 2198 2199 2200 2201 2202 2203 2204 2205 2206 2207 2208 2209 2210 2211 2212 2213 2214 2215 2216 2217 2218 2219 2220 2221 2222 2223 2224 2225 2226 2227 2228 2229 2230 2231 2232 2233 2234 2235 2236 2237 2238 2239 2240 2241 2242 2243 2244 2245 2246 2247 2248 2249 2250 2251 2252 2253 2254 2255 2256 2257 2258 2259 2260 2261 2262 2263 2264 2265 2266 2267 2268 2269 2270 2271 2272 2273 2274 2275 2276 2277 2278 2279 2280 2281 2282 2283 2284 2285 2286 2287 2288 2289 2290 2291 2292 2293 2294 2295 2296 2297 2298 2299 2300 2301 2302 2303 2304 2305 2306 2307 2308 2309 2310 2311 2312 2313 2314 2315 2316 2317 2318 2319 2320 2321 2322 2323 2324 2325 2326 2327 2328 2329 2330 2331 2332 2333 2334 2335 2336 2337 2338 2339 2340 2341 2342 2343 2344 2345 2346 2347 2348 2349 2350 2351 2352 2353 2354 2355 2356 2357 2358 2359 2360 2361 2362 2363 2364 2365 2366 2367 2368 2369 2370 2371 2372 2373 2374 2375 2376 2377 2378 2379 2380 2381 2382 2383 2384 2385 2386 2387 2388 2389 2390 2391 2392 2393 2394 2395 2396 2397 2398 2399 2400 2401 2402 2403 2404 2405 2406 2407 2408 2409 2410 2411 2412 2413 2414 2415 2416 2417 2418 2419 2420 2421 2422 2423 2424 2425 2426 2427 2428 2429 2430 2431 2432 2433 2434 2435 2436 2437 2438 2439 2440 2441 2442 2443 2444 2445 2446 2447 2448 2449 2450 2451 2452 2453 2454 2455 2456 2457 2458 2459 2460 2461 2462 2463 2464 2465 2466 2467 2468 2469 2470 2471 2472 2473 2474 2475 2476 2477 2478 2479 2480 2481 2482 2483 2484 2485 2486 2487 2488 2489 2490 2491 2492 2493 2494 2495 2496 2497 2498 2499 2500 2501 2502 2503 2504 2505 2506 2507 2508 2509 2510 2511 2512 2513 2514 2515 2516 2517 2518 2519 2520 2521 2522 2523 2524 2525 2526 2527 2528 2529 2530 2531 2532 2533 2534 2535 2536 2537 2538 2539 2540 2541 2542 2543 2544 2545 2546 2547 2548 2549 2550 2551 2552 2553 2554 2555 2556 2557 2558 2559 2560 2561 2562 2563 2564 2565 2566 2567 2568 2569 2570 2571 2572 2573 2574 2575 2576 2577 2578 2579 2580 2581 2582 2583 2584 2585 2586 2587 2588 2589 2590 2591 2592 2593 2594 2595 2596 2597 2598 2599 2600 2601 2602 2603 2604 2605 2606 2607 2608 2609 2610 2611 2612 2613 2614 2615 2616 2617 2618 2619 2620 2621 2622 2623 2624 2625 2626 2627 2628 2629 2630 2631 2632 2633 2634 2635 2636 2637 2638 2639 2640 2641 2642 2643 2644 2645 2646 2647 2648 2649 2650 2651 2652 2653 2654 2655 2656 2657 2658 2659 266